



ADEAR

Alzheimer's Disease Education & Referral Center

A Service of the [National Institute on Aging](#)



Enter Search

Search ADEAR

Search ADEAR and NIA

[Home](#) | [About Us](#) | [Site Map](#) | [Sign Up for Alerts](#) | [Contact Us](#) | [Help Viewing Site](#)

Alzheimer's Disease

[General Info](#)
[Causes](#)
[Symptoms](#)
[Diagnosis](#)
[Treatment](#)

Caregiving

[Home/Family](#)
[Professional](#)

Health Info

[News](#)

[Clinical Trials](#)

[AD Research Centers](#)

Publications

[Browse/Search](#)
[Newsletter](#)

Resources

[Literature Search \(CHID\)](#)

[Media Room](#)

[Conference Calendar](#)

Understanding Stages and Symptoms of Alzheimer's Disease*

Alzheimer's disease develops slowly and causes changes in the brain long before there are obvious changes in a person's memory, thinking, use of words or behavior. Stages and changes the person will go through are outlined below.

- [Common Changes in Mild AD](#)
- [Common Changes in Moderate AD](#)
- [Common Changes in Severe AD](#)

Common Changes in Mild AD

- Loses spark or zest for life - does not start anything.
- Loses recent memory without a change in appearance or casual conversation.
- Loses judgment about money.
- Has difficulty with new learning and making new memories.
- Has trouble finding words - may substitute or make up words that sound like or mean something like the forgotten word.
- May stop talking to avoid making mistakes.
- Has shorter attention span and less motivation to stay with an activity.
- Easily loses way going to familiar places.
- Resists change or new things.
- Has trouble organizing and thinking logically.
- Asks repetitive questions.
- Withdraws, loses interest, is irritable, not as sensitive to others' feelings, uncharacteristically angry when frustrated or tired.
- Won't make decisions. For example, when asked what she wants to eat, says "I'll have what she is having."
- Takes longer to do routine chores and becomes upset if rushed or if something unexpected happens.
- Forgets to pay, pays too much, or forgets how to pay - may hand the checkout person a wallet instead of the correct amount of money.
- Forgets to eat, eats only one kind of food, or eats constantly.
- Loses or misplaces things by hiding them in odd places or forgets where things go, such as putting clothes in the dishwasher.
- Constantly checks, searches or hoards things of no value.

Common Changes in Moderate AD

- Changes in behavior, concern for appearance, hygiene, and sleep become more noticeable.
- Mixes up identity of people, such as thinking a son is a brother or that a wife is a stranger.
- Poor judgment creates safety issues when left alone - may wander and risk exposure, poisoning, falls, self-neglect or exploitation.
- Has trouble recognizing familiar people and own objects; may take things that belong to others.
- Continuously repeats stories, favorite words, statements, or motions like tearing tissues.
- Has restless, repetitive movements in late afternoon or evening, such as pacing, trying doorknobs, fingering draperies.
- Cannot organize thoughts or follow logical explanations.
- Has trouble following written notes or completing tasks.
- Makes up stories to fill in gaps in memory. For example might say, "Mama will come for me when she gets off work."
- May be able to read but cannot formulate the correct response to a written request.
- May accuse, threaten, curse, fidget or behave inappropriately, such as kicking, hitting, biting, screaming or grabbing.
- May become sloppy or forget manners.
- May see, hear, smell, or taste things that are not there.
- May accuse spouse of an affair or family members of stealing.
- Naps frequently or awakens at night believing it is time to go to work.
- Has more difficulty positioning the body to use the toilet or sit in a chair.
- May think mirror image is following him or television story is happening to her.
- Needs help finding the toilet, using the shower, remembering to drink, and dressing for the weather or occasion.
- Exhibits inappropriate sexual behavior, such as mistaking another individual for a spouse. Forgets what is private behavior, and may disrobe or masturbate in public.

Common Changes in Severe AD

- Doesn't recognize self or close family.
- Speaks in gibberish, is mute, or is difficult to understand.
- May refuse to eat, chokes, or forgets to swallow.
- May repetitively cry out, pat or touch everything.
- Loses control of bowel and bladder.
- Loses weight and skin becomes thin and tears easily.
- May look uncomfortable or cry out when transferred or touched.
- Forgets how to walk or is too unsteady or weak to stand alone.
- May have seizures, frequent infections, falls.
- May groan, scream or mumble loudly.

Sleeps more.

- Needs total assistance for all activities of daily living.

*Adapted from *Caring for People with Alzheimer's Disease: A Manual for Facility Staff* (2nd edition), by Lisa P. Gwyther, 2001. Published by the [American Health Care Association](#) (1201 L Street, NW, Washington, DC 20005) and the [Alzheimer's Association](#) (919 N. Michigan Ave., Suite 1100, Chicago, IL 60611).

[Back to Top](#)

[Privacy Statement](#) | [Web Accessibility Statement](#) | [Customer Service Standards](#) | [Contact Us](#)